

SWEETLIX BLOAT GUARD (11453)

Feed Attributes

- Medicated with poloxalene for the prevention of legume (alfalfa, clover) and wheat pasture bloat in cattle.
- To be used as a top dressing on individual rations of ground feed.
- Start use 2 or 3 days prior to exposure to bloat-producing conditions and continue use as long as cattle are exposed to bloat-producing conditions.
- Convenient measuring scoop included.

Feeding and Management

1. To be fed at the rate of 0.8 oz. of block per 100 lb. of body weight per day. Example: 8 oz. (0.5 lb.) of block for a 1000 lb. animal daily.
2. For adequate protection it is essential that each animal consume the total recommended dosage of SWEETLIX® Bloat Guard® Block daily.
3. Omit all salt from feed. Do not feed free choice sources of salt or mineral containing salt in any form. These blocks are an adequate source of salt.
4. Begin feeding blocks continuously. Feed blocks and full feed dry non-legume hay at least 48 hours prior to legume (alfalfa, clover) consumption. Repeat when block consumption is interrupted to maintain control.
5. Provide at least 1 block for each 5 head of cattle; add 1 additional block when each block has been half-consumed.
6. The location of the block is extremely important for adequate consumption. Place blocks where cattle congregate (watering, grazing and loafing areas) in order to limit the distance an animal must travel to have ready access to a block.
7. Controlled grazing practices (Strip Grazing) are recommended in order to limit the distance an animal must travel to have ready access to a block.
8. In some instances it may be necessary to confine cattle in a dry lot with additional SWEETLIX Bloat Guard Blocks for a period each day.

WARNING

Access to, and intake of blocks, may be limited during or directly after heavy rain or dew, and after frost or change of pasture. Variation in cattle routine may change their feeding and watering habits. Relocate blocks immediately as necessary. Special caution should be taken. Water high in salt and alkaline soils limit consumption and performance.

CAUTION

USE AS DIRECTED. Consumption of selenium should not exceed 3 mg per head daily.

Guaranteed Analysis

Crude Protein	(Min)	4.0%
Crude Fat	(Min)	0.5%
Crude Fiber	(Max)	12.5%
Salt	(Min)	19.5%
Salt	(Max)	23.0%
Potassium	(Min)	1.8%
Iodine	(Min)	43 ppm
Selenium	(Min)	13 ppm
Vitamin A	(Min)	None Added

List of Ingredients

Cane Molasses, Soybean Hulls, Salt, Hydrolyzed Animal Fat, Manganous Oxide, Zinc Oxide, Ferrous Carbonate, Copper Sulfate, Potassium Iodide, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydriodide and Calcium Carbonate.

33# SBG