

Western Milling House Calf Feed w/Rumensin

Increased rate of weight gain; For prevention and control of coccidiosis caused by Eimeria bovis and Eimeria zuernii.

ACTIVE DRUG INGREDIENT
 Monensin (as Monensin sodium) 44.4 g/ton

Guaranteed Analysis			
Crude protein	(Min)	17.0	%
Crude fat	(Min)	3.1	%
Crude fiber	(Max)	7.0	%
Ash	(Max)	10.6	%
Phosphorus	(Min)	0.4	%
Calcium	(Min)	0.9	%
Calcium	(Max)	1.7	%
Sodium	(Max)	0.3	%
Selenium - added	(Min)	0.6	ppm
Selenium - added	(Max)	1.1	ppm

List of Ingredients

Rolled Corn, Canola Meal, Soybean Meal, Molasses, Rolled Barley, Wheat Millrun, Recleaned oats (fancy), Calcium Carbonate, Alfalfa meal, Animal/Vegetable liquid Fat, Salt, Almond Shell Ground (Not more than 0.6%), Dicalcium Phosphate, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E supplement, Manganese Sulfate, Zinc Sulfate, Sodium Selenite, Copper Sulfate, Potassium Chloride, Magnesium Oxide, Cobalt Sulfate, Ethylenediamine Dihydroiodide, Vegetable Oil, Monensin, Choline Chloride, D-Alpha Tocopheroyl Acetate, Rice Bran, Biotin Supplement, Thiamine Mononitrate, Riboflavin, Folic Acid, Niacin Supplement, Vitamin B12 Supplement.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION: Follow label directions - this feed contains added selenium. Feeding selenium at levels in the total diet in excess of 0.3 ppm is prohibited. This feed contains 1.1 ppm selenium. **A)** Mix this ration with other feeds not to exceed 0.3 ppm of selenium per ton. Do not exceed more than 550 pounds per ton in total mixed ration as sole source of selenium. Consult your nutritionist for proper feeding instructions.

One LB of feed will provide 22.2 mg of Monensin. 9.0 lb will provide the maximum level of 200 mg.

Pasture cattle (slaughter, stocker, and feeder dairy & beef replacement heifers). For increased rate of weight gain. Feed at the rate of not less than 50mg nor more than 200 mg per head per day in not less than one pound of feed. During the first 5 days, cattle should receive no more than 100 mg per day contained in not less than 1 pound of feed. **DO NOT SELF FEED.** For the prevention and control of coccidiosis due to eimeria bovis & elmeria zuernii. Feed at a rate to provide 0.14 TO 0.42 mg per pound body weight per day depending upon severity of challenge up to a maximum 200 mg per head per day. During the first 5 days, cattle should receive no more than 100 mg per day contained in not less than 1 pound of feed.

83985 50#



Promises Made. Promises Kept.