

K CHICK STARTER MASH

As an aid in prevention of coccidiosis caused by Eimeria tenella only.

ACTIVE DRUG INGREDIENT

Amprolium0.0125%

Feeding Instructions

Feed as the sole ration to chicks from day-old until pullets are 6 weeks old. Always provide fresh, clean water.

WARNING

Withdraw 5 days before slaughter. Use as a sole source of Amprolium.

Special Considerations

Do not use Amprolium in feeds containing bentonite.

Guaranteed Analysis

CRUDE PROTEIN (Min)	21.0%
CRUDE FAT (Min)	2.3%
CRUDE FIBER (Max)	5.4%
ASH (Max)	9.7%
PHOSPHORUS (Min)	0.6%
CALCIUM (Min)	0.8%
CALCIUM (Max)	1.5%

List of Ingredients

Ground Corn, Soybean Meal, Wheat Millrun, Rice Bran, Calcium Carbonate, Dicalcium Phosphate, Molasses, Amprolium 2.5%, Salt, Choline Chloride, Iron Sulfate, Manganous Oxide, Zinc Oxide, Ethoxyquin, Niacin, Vitamin E Supplement, Sodium Selenite, Copper Sulfate, D-Calcium Pantothenate, Ethylenediamine Dihydroiodide, Biotin, Menadione, Pyridoxine Hydrochloride, Riboflavin, Vitamin B12 Supplement, Folic Acid, Vitamin A Supplement, Vitamin D3 Supplement, Methionine Liquid.

50# K53605