

Showstring Beef Grower/Finisher

Improved feed efficiency; for reduction of incidence of liver abscesses caused by *Fusobacterium necrophorum* and *Arcanobacterium (Actinomyces) pyogenes*.

ACTIVE DRUG INGREDIENTS

Monensin 30 g/ton Tylosin 10 g/ton

Guaranteed Analysis

Crude protein*	(Min)	13.0	%
* (INCLUDES NOT MORE THAN 1.5 % EQUIVALENT CRUDE PROTEIN FROM NON-PROTEIN NITROGEN)			
Crude fat	(Min)	4.5	%
Crude fiber	(Max)	11.2	%
Ash	(Max)	9.3	%
Phosphorus	(Min)	0.3	%
Calcium	(Min)	0.5	%
Calcium	(Max)	1.0	%
Sodium	(Max)	0.6	%

List of Ingredients

Rolled Corn, Soybean Meal, Cottonseed Hulls (Not more than 11.4%), Molasses, Rolled Barley, Beet Pulp Shreds, Recleaned oats (fancy), Cottonseed Whole White, Animal/Vegetable liquid Fat, Wheat Millrun, Sodium Sesquicarbonate, Calcium Carbonate, Dicalcium Phosphate, Corn Distillers, Linseed Meal Expeller, Urea, Sodium Silico Aluminate, Vegetable Oil, Salt, Whey Dried, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E supplement, Brewers Dried Yeast, Dried *Saccharomyces cerevisiae* fermentation solubles, Brewers Yeast Dehydrated, Kelp (Seaweed Meal), Zinc Methionine, Manganese Methionine, Copper Lysine, Cobalt Glucoheptoate, Anise flavor dry, Magnesium Oxide, Monensin, Tylosin Phosphate, Sulfur, Selenium Yeast, Brewers Dried Yeast, Propionic Acid, Ammonium Hydroxide, Vexite, Bentonite, Artificial flavoring, *Saccharomyces cerevisiae* yeast, Cane Molasses, Diastatic Malt, Corn Syrup, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Ferrous Sulfate, Niacin, Riboflavin Supplement, Calcium Pantothenate, Biotin, Choline Chloride, Sodium Selenite, Thiamine Mononitrate, Vitamin B-12 Supplement, Folic Acid, Calcium Iodate, Rice Bran, Cobalt sulfate.. **CAUTION:** Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Do not feed to lactating dairy cattle. Consumption by unapproved species may result in toxic reactions. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Feeding undiluted or mixing errors resulting in high concentrations of Monensin could be fatal to cattle. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. **LIMITATIONS:** Do not use in any feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite. **WARNING: EXCESSIVE CONSUMPTION MAY RESULT IN ADVERSE TOXIC REACTION - USE ONLY AS DIRECTED CAUTION:** This feed contains a source of non-protein nitrogen. Provide adequate drinking water and do not use with other feed containing non-protein nitrogen. Caution should be exercised with, starved, stressed or debilitated animals. All manufacturer's directions for use must be followed carefully. **FOR RUMINATES ONLY** Feed Grain concentrate at 2-3% of body weight along with roughage. Provide free choice salt and fresh clean water at all time.

500015 50#



Promises Made. Promises Kept.

5/14/2013