

# Showstring Creep/Developer

Improved feed efficiency; for the reduction of incidence of the liver abscesses caused by Fusobacterium Necrophorum and Actinomyces (Actinomyces) Pyogenes.

Monensin ..... 30 g/ton  
Tylosin ..... 10 g/ton

## Guaranteed Analysis

Crude protein\* (Min) 14.0 %

\* (INCLUDES NOT MORE THAN 1.6 % EQUIVALENT CRUDE PROTEIN FROM NON-PROTEIN NITROGEN)

Crude fat	(Min)	3.5 %
Crude fiber	(Max)	10.0 %
Ash	(Max)	8.5 %
Phosphorus	(Min)	0.4 %
Calcium	(Min)	0.6 %
Calcium	(Max)	1.2 %
Sodium	(Max)	0.3 %

## List of Ingredients

Recleaned oats (fancy), Rolled Corn, Soybean Meal, Beet Pulp Shreds, Molasses, Alfalfa meal, Cottonseed Whole White, Wheat Millrun, Calcium Carbonate, Dicalcium Phosphate, Corn Distillers, Linseed Meal Expeller, Urea, Sodium Silico Aluminate, Vegetable Oil, Salt, Sodium Sesquicarbonate, Whey Dried, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E supplement, Brewers Dried Yeast, Dried Saccharomyces cerevisiae fermentation solubles, Brewers Yeast Dehydrated, Kelp (Seaweed Meal), Zinc Methionine, Manganese Methionine, Copper Lysine, Cobalt Glucoheptoate, Anise flavor dry, Magnesium Oxide, Monensin, Tylosin Phosphate, Sulfur, Selenium Yeast, Brewers Dried Yeast, Saccharomyces cerevisiae yeast, Cane Molasses, Diastatic Malt, Corn Syrup, Propionic Acid, Ammonium Hydroxide, Verxite, Bentonite, Artificial flavoring, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Ferrous Sulfate, Niacin, Riboflavin Supplement, Calcium Pantothenate, Biotin, Choline Chloride, Sodium Selenite, Thiamine Mononitrate, Vitamin B-12 Supplement, Folic Acid, Calcium Iodate, Rice Bran, Cobalt sulfate.. **CAUTION:** Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Do not feed to lactating dairy cattle. Consumption by unapproved species may result in toxic reactions. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Feeding undiluted or mixing errors resulting in high concentrations of Monensin could be fatal to cattle. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. **LIMITATIONS:** Do not use in any feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite. **WARNING: EXCESSIVE CONSUMPTION MAY RESULT IN ADVERSE TOXIC REACTION - USE ONLY AS DIRECTED CAUTION:** This feed contains a source of non-protein nitrogen. Provide adequate drinking water and do not use with other feed containing non-protein nitrogen. Caution should be exercised with, starved, stressed or debilitated animals. All manufacturer's directions for use must be followed carefully. **FOR RUMINATES ONLY** Feed Grain concentrate at 2-3% of body weight along with roughage. Provide free choice salt and fresh clean water at all time.

500085 50#



Promises Made. Promises Kept.