

SHOWSTRING HEIFER MIX

Improved feed efficiency; for reduction of incidence of liver abscesses caused by *Fusobacterium necrophorum* and *Actinomyces (Corynebacterium) pyogenes*.

ACTIVE DRUG INGREDIENTS

Monensin	26 g/ton
Tylosin	8 g/ton

Guaranteed Analysis

Crude protein*	(Min)	13.5 %
* (INCLUDES NOT MORE THAN 1.0 % EQUIVALENT CRUDE PROTEIN FROM NON-PROTEIN NITROGEN)		
Crude fat	(Min)	2.2 %
Crude fiber	(Max)	16.7 %
Ash	(Max)	7.8 %
Copper - added	(Min)	24 ppm
Copper - added	(Max)	29 ppm

List of Ingredients

Beet Pulp Shreds with Molasses, Recleaned oats (fancy), Rolled Barley, Cottonseed Hulls (Not more than 13.0%), Soy hull pellets (Not more than 12.0%), Soybean Meal, Molasses, Canola Meal, Wheat Millrun, Calcium Carbonate, Ground Corn, Dicalcium Phosphate, Corn Distillers, Almond Shell Ground (Not more than 0.5%), Linseed Meal Expeller, Urea, Sodium Silico Aluminate, Salt, Vegetable Oil, Sodium Sesquicarbonate, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, dl-Alpha Tocopheryl Acetate, Selenium Yeast, Ferrous Sulfate, Niacin, Riboflavin Supplement, Vitamin A Supplement, Calcium Pantothenate, Biotin, Choline Chloride, Sodium Selenite, Thiamine Mononitrate, Vitamin D3 Supplement, Vitamin B-12 Supplement, Folic Acid, Calcium Iodate, Rice Bran, Cobalt sulfate., Whey Dried, *Saccharomyces cerevisiae* yeast, Cane Molasses, Diastatic Malt, Corn Syrup, Fish Meal, Brewers Dried Yeast, Dried *Saccharomyces cerevisiae* fermentation solubles, Brewers Yeast Dehydrated, Zinc Methionine, Manganese Methionine, Copper Lysine, Cobalt Glucoheptoate, Anise flavor dry, Kelp (Seaweed Meal), Alfalfa Meal, Iron Sulfate, Manganous Oxide, Zinc Oxide, Ethoxyquin, Vitamin E Supplement, D-Calcium Pantothenate, Ethylenediamine Dihydroiodide, Menadione, Pyridoxine Hydrochloride, Riboflavin, Vitamin B12 Supplement, Propionic Acid, Ammonium Hydroxide, Verxite, Bentonite, Artificial flavoring, Monensin, Magnesium Oxide, Tylosin Phosphate, Iron Oxide, Sulfur, Brewers Dried Yeast, Methionine Liquid, Lysine, Red Color Dye.

CAUTION: Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Feeding undiluted or mixing errors resulting in high concentrations of Monensin could be fatal to cattle. Must be thoroughly mixed in feeds before use; do not feed undiluted.

LIMITATIONS: Feed only to cattle being fed in confinement for slaughter. Feed continuously at the rate of 8.2 to 10.2 kilograms (18 to 22.5 pounds) of Type C medicated feed per head per day to supply 240 milligrams of monensin and 90 milligrams of tylosin per head per day; as monensin sodium; as tylosin phosphate. Safe use in unapproved species and breeding cattle has not been established.

CAUTION: This feed contains added levels of copper do not feed to sheep.

WARNING: EXCESSIVE CONSUMPTION MAY RESULT IN ADVERSE TOXIC REACTION - USE ONLY AS DIRECTED

CAUTION: This feed contains a source of non-protein nitrogen. Provide adequate drinking water and do not use with other feed containing non-protein nitrogen. Caution should be exercised with, starved, stressed or debilitated animals. All manufacturer's directions for use must be followed carefully. FOR RUMINATES ONLY

500115 50#



Promises Made. Promises Kept.