

1448



0 0 0 0 1 6 0 4 9

SHOW PIG 806 PRESTARTER 23% 50#

This feed is designed to be fed to starting swine from 3 to 9 weeks of age. **MEDICATED:** Control of swine dysentery associated with Brachyspira (formerly Serpulina or Treponema) hydodysenteriae susceptible to Tiamulin. Treatment of bacterial enteritis caused by Escherichia coli and Salmonella choleraesuis and bacterial pneumonia caused by Pasteurella multocida susceptible to Chlortetracycline.

ACTIVE DRUG INGREDIENTS

Tiamulin hydrogen fumarate 35 g/ton Chlortetracycline 400 g/ton

GUARANTEED ANALYSIS

Crude Protein	Min.	23.0	%		
Lysine	Min.	1.6	%		
Crude Fat	Min.	6.5	%		
Crude Fiber	Max.	4.5	%		
Calcium	Min.	0.8	%	Max.	1.1 %
Phosphorus	Min.	0.7	%		
Sodium	Min.	0.1	%	Max.	0.5 %
Copper	Min.	240	PPM		
Zinc	Min.	3300	PPM		

INGREDIENTS: Dehulled Soybean Meal, Ground Corn, Dried Whey, Oat Groats, Soy Hulls (Not more than 4.5%), Rice Bran, Fish Meal, Animal Fat, Vegetable Oil, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Choline Chloride, Iron Sulfate, Zinc Sulfate, Manganous Oxide, Vitamin E Supplement, Vitamin B-12 Supplement, Copper Sulfate, Niacin, Calcium Pantothenate, Vitamin A Supplement, Menadione Sodium Bisulfite Complex, Riboflavin, Folic Acid, Vitamin D3 Supplement, Biotin, Pyridoxine Hydrochloride, Ethylenediamine Dihydriodide, Thiamine Mononitrate, Sodium Selenite, L-Lysine Monohydrochloride, Zinc Oxide, Chlortetracycline, Calcium Carbonate, DL-Methionine Hydroxy Analogue, Brewers Dried Yeast, Dried Saccharomyces Cerevisiae Fermentation Solubles, Diatomaceous Earth, Natural and Artificial Flavors added, Propionic Acid (a preservative), L-Carnitine, Chromium Tripicolinate, Zinc Proteinate, Manganese Proteinate, Copper Proteinate, Yeast, Fermentation Solubles, Yucca Schidigeria

FEEDING DIRECTIONS: Feed continuously as sole ration for 14 days. Not for use in swine weighing more than 250 lb. Feed approximately 400 g/ton of Chlortetracycline, varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight per day. Provide plenty of fresh, clean water at all times

WARNING: Withdraw 2 days before slaughter. CAUTION: Do not feed to animals other than swine. Use as the only source of Tiamulin and Chlortetracycline. The effects of Tiamulin on swine reproductive performance, pregnancy, and lactation have not been determined. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Monensin, Lasalocid, Narasin, Salinomycin, and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medication. **SPECIAL CONSIDERATION:** Protect medicated feeds from excessive temperatures. Do not store at warm temperatures, 104F (40C), for more than 7 days.

MANUFACTURED BY:

O.H. KRUSE GRAIN AND MILLING


31120 West Street

GOSHEN, CA 93227 PH 877-455-6313

274125 B 02/18/16

Net Wt. 50 lbs (22.68 kg)

MEDICATED



2 7 4 1 2 5