

1609



0 0 0 0 1 6 0 4 9

# SHOW PIG 812 STARTER 21% 50#

This feed is designed to be fed to starting swine between 3 to 9 weeks of age. **MEDICATED** -- Control of swine dysentery associated with Brachyspira (formerly Serpulina or Treponema) hyodysenteriae susceptible to Tiamulin. Treatment of bacterial enteritis caused by Escherichia coli and Salmonella choleraesuis and bacterial pneumonia caused by Pasteurella multocida susceptible to Chlortetracycline.

### ACTIVE DRUG INGREDIENTS

Tiamulin hydrogen fumarate .... 35 g/ton      Chlortetracycline ..... 400 g/ton

### GUARANTEED ANALYSIS

Crude Protein	Min.	21.0	%		
Lysine	Min.	1.5	%		
Crude Fat	Min.	6.0	%		
Crude Fiber	Max.	5.0	%		
Calcium	Min.	0.7	%	Max.	1.2 %
Phosphorus	Min.	0.7	%		
Sodium	Min.	0.1	%	Max.	0.5 %
Copper	Min.	240	PPM		
Zinc	Min.	3200	PPM		

**INGREDIENTS:** Ground Corn, Dehulled Soybean Meal, Dried Whey, Flaked Oat Groats, Soy Hulls (Not more than 4.5%), Rice Bran, Blood Meal (Ring Dried), Animal Fat and Vegetable Oil, Monocalcium Phosphate, Dicalcium Phosphate, Calcium Carbonate, Salt, Choline Chloride, Iron Sulfate, Zinc Sulfate, Manganous Oxide, Vitamin E Supplement, Vitamin B-12 Supplement, Copper Sulfate, Niacin, Calcium Pantothenate, Vitamin A Supplement, Menadione Sodium Bisulfite Complex, Riboflavin, Folic Acid, Vitamin D3 Supplement, Biotin, Pyridoxine Hydrochloride, Ethylenediamine Dihydriodide, Thiamine Mononitrate, Sodium Selenite, Zinc Oxide, L-Lysine Monohydrochloride, Brewers Dried Yeast, Dried Saccharomyces Cerevisiae Fermentation Solubles, DL-Methionine Hydroxy Analogue, Natural and Artificial Flavors added, Diatomaceous Earth, Propionic Acid (a preservative), Zinc Proteinate, Manganese Proteinate, Copper Proteinate, Yeast, Fermentation Solubles, Dried Yucca Schidigera Powder.

**FEEDING DIRECTIONS:** Each pound of this medicated feed contains 200 mg of Chlortetracycline and 17.5 mg Tiamulin. Feed approximately 400 g/ton of Chlortetracycline, varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight per day - approximately 5 lbs of feed per every 100 lbs bodyweight. Feed for not more than 14 days. Provide plenty of fresh, clean water at all times.

**CAUTION:** Do not feed to animals other than swine. **Withdraw 2 days before slaughter.**

**SPECIAL CONSIDERATON:** Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Monensin, Lasalocid, Narasin, Salinomycin, and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use. The effects of Tiamulin on swine reproductive performance, pregnancy, and lactation have not been determined. Protect medicated feeds from excessive temperatures. Do not store at warm temperatures, 104°F (40°C), for more than 7 days.

MANUFACTURED BY:

**O.H. KRUSE GRAIN AND MILLING**


31120 West Street

GOSHEN, CA 93227 PH 877-455-6313

**274145 O**      02/18/16

Net Wt. 50 lbs (22.68 kg)

**MEDICATED**



2 7 4 1 4 5