

1711



0 0 0 0 1 5 3 4 1



SUNGLO® STARTER-803 MEDICATED

For pigs weighing 18 to 75 pounds bodyweight. **MEDICATED** -- Control of swine dysentery associated with Brachyspira (formerly Serpulina or Treponema) hyodysenteriae susceptible to Tiamulin. Treatment of bacterial enteritis caused by Escherichia coli and Salmonella choleraesuis and bacterial pneumonia caused by Pasteurella multocida susceptible to Chlortetracycline.

ACTIVE DRUG INGREDIENTS

Tiamulin hydrogen fumarate 35 g/ton Chlortetracycline 400 g/ton

GUARANTEED ANALYSIS

Crude Protein	Min.	19.0	%		
Lysine	Min.	1.44	%		
Crude Fat	Min.	5.5	%		
Crude Fiber	Max.	3.0	%		
Calcium	Min.	0.67	%	Max.	1.17 %
Phosphorus	Min.	0.66	%		
Salt	Min.	0.14	%	Max.	0.35 %
Sodium	Min.	0.10	%	Max.	0.50 %
Selenium	Min.	0.30	PPM		
Zinc	Min.	2900	PPM		

INGREDIENTS: Ground Corn, Dehulled Soybean Meal, Wheat Middlings, Dried Whey, Flaked Oat Groats, Animal Fat and Vegetable Oil, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Corn Gluten Meal, Fish Meal, Whey Permeate, Bovine Plasma, Egg Product, L-Lysine, Calcium Carbonate, Zinc Oxide, BHT, Benzoic Acid, Citric Acid, Sorbic Acid, Sodium Benzoate, Formic Acid, Lactic Acid, Fumaric Acid, Sodium Hydroxide, Choline Chloride, DL-Methionine, L-Threonine, Lignin Sulfonate, Copper Sulfate, Ferrous Sulfate, Manganous Oxide, Zinc Sulfate, Calcium Iodate, Sodium Selenite, Saccharomyces cerevisiae, Potassium Chloride, Chromium Tripicolinate, Vitamin E Supplement, d-Calcium Pantothenate, Niacin Supplement, Menadione Sodium Bisulfite Complex, Riboflavin Supplement, Vitamin D3 Supplement, Biotin, Vitamin B12 Supplement, Pyridoxine Hydrochloride, Vitamin A Acetate, Thiamine Mononitrate, Folic Acid, Saccharin Sodium, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Glucoheptonate, Dried Bacillus licheniformis Fermentation Product, Dried Bacillus subtilis Fermentation Product, Yucca Schidigera Extract, L-Tryptophan, Natural and Artificial Flavor

FEEDING DIRECTIONS: Feed as a sole ration to pigs weighing 18 to 75 pounds. Feed continuously as sole ration for 14 days.

WARNING: Withdraw 2 days before slaughter. **CAUTION:** Do not feed undiluted. Do not feed to animals other than swine. The effects of Tiamulin on swine reproductive performance, pregnancy and lactation have not been determined. Use as the only source of Tiamulin and Chlortetracycline. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g. Monensin, Lasalocid, Narasin, Salinomycin and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of mediated feeds. **SPECIAL CONSIDERATIONS:** Protect medicated feeds from excessive temperatures. Do not store at warm temperatures, 104°F (40°C), for more than 7 days. Sunglo® is a registered trademark of Provimi North America, Inc.

MANUFACTURED BY:

O.H. KRUSE GRAIN AND MILLING


31120 West Street

GOSHEN, CA 93227 PH 877-455-6313

73205 P 12/07/15

Net Wt. 50 lbs (22.68 kg)

MEDICATED



7 3 2 0 5