

SWINE CELL-RATE 17% GRW/FIN PLT

For control of swine dysentery caused by *Brachyspira hyodysenteriae*; and for control of porcine proliferative enteropathies (ileitis) associated with *Lawsonia intracellularis*.

ACTIVE DRUG INGREDIENT

Tylosin.....100 g/ton

Feeding Directions

Provide clean fresh water at all times.

Notice

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

100g/ton for at least 3 weeks, followed by 40g/ton until swine reach market weight.

100g/ton maintaining weight gains & feed efficiency in the presence of atrophic rhinitis.

100g/ton for 21 days as sole ration – prevention and/or porcine proliferative enteropathies associated with *Lawsonia intracellularis*.

Guaranteed Analysis

CRUDE PROTEIN (Min)	17.0%
CRUDE FAT (Min)	5.3%
CRUDE FIBER (Max)	4.8%
ASH (Max)	8.5%

List of Ingredients

Ground Corn, Soybean Meal, Whey Dried, Wheat Millrun, Animal Fat/Vegetable liquid fat, Oat Groats, Fish Meal, Dicalcium Phosphate, Calcium Carbonate, Salt, Choline Chloride, Iron Sulfate, Zinc Sulfate, Manganous Oxide, Vitamin E Supplement, Vitamin B-12 Supplement, Copper Sulfate, Niacin, Calcium Pantothenate, Vitamin A Supplement, Menadione Sodium Bisulfite Complex, Riboflavin, Folic Acid, Vitamin D3 Supplement, Biotin, Pyridoxine Hydrochloride, Ethylene Diamine Dihydriodide, Thiamine Mononitrate, Sodium Selenite, Lysine, Tylosin Phosphate, Flavor, Brewers Dried Yeast, Dried *Saccharomyces Cerevisiae* Fermentation Solubles, Yeast Cell Rate, Methionine Liquid, Zinc Oxide, Zinc Proteinate, Manganese Proteinate, Copper Proteinate, Yeast, Fermentation Solu.

50# 228455

MEDICATED

