

# SWINE CELL-RATE 19% STR/GRW PLT

For control of Swine Dysentery and the control of Porcine Proliferative Enteropathy (Ileitis) caused by Lawsonia Intracellularis. Feed as the sole ration.

## ACTIVE DRUG INGREDIENT

Lincomycin ..... 40 g/ton

### Feeding Directions

Provide clean fresh water at all times.

### LIMITATIONS FOR USE

For use in animals or on premises with a history of swine dysentery, but where symptoms have not yet occurred.

### CAUTION

Occasionally, swine fed Lincomycin may, within the first 2 days after the onset of treatment, develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within 5-8 days without discontinuing the Lincomycin treatment. Do not allow rabbits, hamsters, guinea pigs, horses or ruminants access to feeds containing Lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Not to be fed to swine that weigh more than 250 lb.

### Guaranteed Analysis

CRUDE PROTEIN (Min)	19.0%
CRUDE FAT (Min)	5.5%
CRUDE FIBER (Max)	4.7%
ASH (Max)	9.5%
PHOSPHORUS (Min)	0.6%
CALCIUM (Min)	0.8%
CALCIUM (Max)	1.4%
SODIUM (Max)	0.3%

### List of Ingredients

Ground Corn, Soybean Meal, Whey Dried, Wheat Millrun, Animal/Vegetable liquid fat, Oat Groats, Fish Meal, Dicalcium Phosphate, Calcium Carbonate, Salt, Choline Chloride, Iron Sulfate, Zinc Sulfate, Manganous Oxide, Vitamin E Supplement, Vitamin B-12 Supplement, Copper Sulfate, Niacin, Calcium Pantothenate, Vitamin A Supplement, Menadione Sodium Bisulfite Complex, Riboflavin, Folic Acid, Vitamin D3 Supplement, Biotin, Pyridoxine Hydrochloride, Ethylene Diamine Dihydriodide, Thiamine Mononitrate, Sodium Selenite, Zinc Oxide, Yeast Cell Rate, Lincomycin, Flavor, Brewers Dried Yeast, Dried Saccharomyces Cerevisiae Fermentation Solubles, Methionine Liquid, Zinc Proteiniate, Manganese Proteiniate, Copper Proteiniate, Yeast, Fermentation Solu, Lysine.

**50# 228465**

**MEDICATED**

